



LAO PEOPLE 'S DEMOCRATIC REPUBLIC
PEACE INDEPENDENCE DEMOCRACY UNITY PROSPERITY

MINISTRY OF HEALTH
FOOD AND DRUG DEPARTMENT
TEL (856 21) 214013, 214014
FAX: (856 21) 214015.

Certificate of Drug Registration

Registration Number:07-LM-10053-26

ALIAMPI-500

Dosage Form	CAPSULE
Active Ingredients	AMPICILLIN TRIHYDRATE 500 MG
Description	Red-White capsule shell and its contents are white or off-white powder, PVC/Alu blister, 10 capsules/blister, 50 blisters/box
Size of packaging unit	500 CAPSULES
Indications	See Annex
Manufacturer	LAOS ALLIANCE PHARMACEUTICAL GROUP COMPANY SAYSETHA DEVELOPMENT ZONE, SAYSETHA DISTRICT, VIENTIANE CAPITAL, LAO PDR

This Certificate is issued to attest that the product described above has met the minimum standards required for drug registration undertaken by the Food and Drug Department in Lao PDR.

This certificate expires, unless previously suspended or revoked, on **23 July 2031**

Note: All boxes and packaging for importations must bear the Lao Registration No. **07-LM-10053-26**

At, Vientiane, **23 July 2026**



ນ.ພຸດທະວັນ ອິນລໍຄຳ



Registration Number:07-LM-10053-26

ALIAMPI-500

ANNEX

INDICATIONS

Ampicillin is a broad-spectrum antibacterial indicated for treatment of commonly occurring bacterial infections where a sensitive organism is suspected or proven?

- Lower respiratory tract-acute and chronic bronchitis, lobar and bronchopneumonia.
- Upper respiratory tract-bacterial pharyngitis, otitis media, chronic bronchial sepsis.
- Genito- urinary tract-acute cystitis, pyelonephritis for sensitive infections, gonorrhoea (in combination with probenecid.) and gynaecological infections.
- Other - skin and soft tissue infections, dental abscess (as an adjunct to surgical management), enteric fever, intra-abdominal sepsis, peritonitis and septicaemia (in combination with an aminoglycoside or metronidazole).

Ampicillin ແມ່ນຢາຕ້ານເຊື້ອທີ່ກວມຄວບຄຸມກວ້າງ (Broad-spectrum antibacterial) ເຊິ່ງໃຊ້ເຂົ້າໃນການປິ່ນປົວພະຍາດຕິດເຊື້ອແບັກທີເຣຍຫຼາຍໆ ໃນກໍລະນີທີ່ສົງໄສ ຫຼື

ຍັງບໍ່ຢືນຢັນວ່າເກີດຈາກເຊື້ອທີ່ຕອບສະໜອງຕໍ່ຢານີ້:

- ລະບົບຫາຍໃຈສ່ວນລຸ່ມ: ຫຼອດລົມອັກເສບກະທັນຫັນ ແລະ ຊ້າເຮືອ, ປອດອັກເສບ.
- ລະບົບຫາຍໃຈສ່ວນເທິງ: ຄໍອັກເສບຈາກເຊື້ອແບັກທີເຣຍ, ຫູສ່ວນກາງອັກເສບ, ຕິດເຊື້ອໃນຫຼອດລົມຊ້າເຮືອ.
- ລະບົບທາງເດີນປັດສະວະ ແລະ ສືບພັນ: ຝຶກຍ່ຽວອັກເສບ, ໝາກໄຂ່ຫຼັງອັກເສບ, ພະຍາດໝອງໃນ (ໃຊ້ຮ່ວມກັບ Probenecid) ແລະ ການຕິດເຊື້ອທາງລະບົບສືບພັນຍິງ.
- ການຕິດເຊື້ອອື່ນໆ: ການຕິດເຊື້ອທາງຜິວໜັງ ແລະ ເນື້ອເຍື່ອອ່ອນ, ຝີແຂ້ວ (ໃຊ້ຄວບຄູ່ກັບການຜ່າຕັດ), ໄຂ້ທໍລະພິດ, ການຕິດເຊື້ອໃນຊ່ອງທ້ອງ, ເຍື່ອບຸຊ່ອງທ້ອງອັກເສບ ແລະ ການຕິດເຊື້ອໃນກະແສເລືອດ (ໃຊ້ຮ່ວມກັບກຸ່ມ Aminoglycoside ຫຼື Metronidazole).